Implementation of LIGO's Triggered Burst Search

Rauha Rahkola (U of Oregon),
representing the
LIGO Scientific Collaboration

This material is based upon work supported in part by National Science Foundation Award PHY-0071058

May 30-31, 2003,
LIGO G030263-00-Z

APS NW Section Meeting
Reed College, Portland, OR

LIGO in Brief





LIGO interferometers (IFOs) measure gravity waves in strain (h) – *the amplitude of perturbation about Minkowski spacetime*

Triggered Burst Search: Motivation



GRB020813 Illustration



Credit: Illustration: CXC/M.Weiss; Spectrum: NASA/CXC/N.Butler et al.

Gravity Wave Bursts (GWBs) are by definition unmodeled!!

Question:

How can we use what we know† (astrophysics) to look for things we don't know (GWBs)?

Answer:

We can use astrophysical "triggers" to point out likely times for GWB events

†or what we think we know

Triggered Burst Search: Overview



- Trigger time distinguishes "on-source" data from "off-source" data for each IFO.
- "On-source" Data: GWB likely
  - GWB arrives before trigger
  - GWB duration is short (~1ms), but exact arrival time is uncertain!
- "Off-source" Data: GWB **not** likely
  - Well past any model-predicted GWBs
  - Close to trigger time to ensure IFO hasn't changed
- How can we use the fact that LIGO has three similar interferometers?

Triggered Burst Search: Technique

1. Filter the data for each IFO
2. Translate source directions to Δt_{GWB}
3. Shift data by Δt_{GWB}
4. FFT & cross-correlate
5. "c.c. statistic" – Σa_{freq} over some frequency band
6. Compare the c.c. statistic for "on-source" data vs. "off-source" data





Sensitivity of the Technique

Expected sensitivity using "off-source" c.c. distribution at a random time over frequencies (200-600Hz):

(95% confidence) ~ $\sqrt{(1.65\sigma)}$ = 5.18E-21 (strain/$\sqrt{Hz}$)

"Off-source" distribution of c.c. statistics
(mean = -0.001313, std. dev. = 1.62932E-41)



Positive GWB signal would compare as an outlier (single trigger case),

OR

as a distribution with +-shifted mean (multiple trigger case)

Current Analysis Issues



Issues which cannot be controlled:

- stability/ sensitivity of the two-IFO system

Variations within trigger sources:

- GWB duration
- GWB source direction/polarization ("antenna pattern")
- GWB source frequencies



Courtesy http://space.mit.edu/hete/bursts/grb030329/

Issues for improving the method:

- Combining single-event results
- Meeting conditions for a multi-trigger analysis
- Dealing with unknown GWB arrival times/durations
- Evaluating tolerance to non-stationarity
- Combining 3+ detectors
- Incorporating current models/classes of models

Summary

- Detecting GWB sources can be optimized using astrophysical triggers
- Cross-correlating data from two similar detectors elicits the coherent GWB signal
- Comparing “on-source” and “off-source” c.c. statistics yields the GW strain associated with a trigger
- Obtaining the associated GW strain can test models of GRB phenomena

LIGO

Possible questions:

- How to deal with differences in IFO response functions
  - Scaling factor proportional to the average calibration difference
- Can you weight the strain with the observed distance to get energy at the source?
  - Possibly
- How can you test specific models of GRBs?
  - Slides 11,12
- What are antenna patterns, and why should they look like that?
  - Slide 13
- What are some models of GRBs?
  - Slide 14
- What are the causes for non-stationary frequencies?
  - Slide 15, but no pictures!
- Why is stationarity/Gaussianity important?
  - Slide 16
- What is the size of burst we injected on slide 3?
  - h_{peak} =9.5E-19
- How sensitive to error boxes for the sources?
  - Depends on the difference in light travel time; approx. 60μsec intervals
- Response time of the IFO to the GW?
  - On the order of μsecs

Interpreting Model-Dependency



- Final result is an estimate of the observable band-limited rms strain (h_{rms}) associated with the trigger
- Apply to an approximate waveform to test models
- Need to account for missing non-stationary frequencies!

Incorporating Model-Dependency

Various Options:

- Divide triggers into groups with similar characteristics
- Use a model-dependent "kernel" in cross-correlation
- Tune to different integration lengths, arrival times

Pictures:

- 4-spectra plot with IFO1, IFO2, kernel, cross-correlation

LIGO's Antenna Patterns

- Polarization-dependent
- Optimally sensitive to GWs propagating normally to detector
- Minimally (but not always zero!) sensitive in the plane of the detector
- Zero sensitivity in the plane of the detector, 45º between the arms

two polarizations: '+' and '×'



unpolarized



GRB Models

Methods to produce GRBs:

- "Exploding Fireball"
- "Black Hole/Magnetic Torus"
- "Cannonball"

GRB progenitors:

- Hypernovae
- Merging of a binary system

Sources of non-stationarity

Figures:

- Violin mode resonances
- Some servo filter response functions

- Mechanical thermal resonances (suspending wire, internal modes of the optics)
- Servo resonances
- 60Hz electronics noise
- Acoustic coupling to the laboratory

Why Gaussianity is Important

- Gaussianity yields predictable statistics
- Gaussianity implies stationarity on a given timescale
- Gaussianity allows several triggers to be combined for better certainty

Gaussianity before and after data conditioning