
Beating the Crab pulsar spin-down and other upper limits: new results from the LIGO S5 known pulsar search


Matthew Pitkin for the LIGO Scientific
Collaboration

`matthew@astro.gla.ac.uk`

LIGO-G070230-00-Z


Targeted pulsar search - Why?

- Rapidly spinning neutron stars provide a potential source of continuous gravitational waves
- To emit gravitational waves they must have some degree of non-axisymmetry e.g. triaxial deformation due to elastic stresses or magnetic fields
- Size of distortions can reveal information about the neutron star equation of state
- Many millisecond and fast young pulsars have very well determined parameters and are generally very stable - good candidates for a targeted search using gravitational detectors!
- The greatly reduced unknown parameter space allows deep, relatively computational inexpensive searches using long time spans of data



Targeted search - method

- Pulsar signal parameterised by: h_0 - gravitational wave amplitude, ι - pulsar orientation, ϕ - gravitational wave phase, and ψ - polarisation angle
- Signal is Doppler modulated by detector and source motion
- Assume that the gravitational wave emission from triaxial neutron star that is tightly coupled and phase locked with the EM emission
 - » Heterodyne time domain data using the known phase evolution of the pulsar
 - » Bayesian parameter estimation of unknown pulsar parameters using data from all interferometers
- Within the LIGO sensitive band ($\nu_{\text{gw}} > 50$ Hz) there are currently 163 known pulsars
- We have rotational and positional parameter information for 97 of these



$$h_0 = \frac{16 \pi^2 G}{c^4} \frac{I \varepsilon \nu_r^2}{d}$$

$$0.95 = \int_{h_0=0}^{h_0^{95}} dh_0 \iiint p(a | \text{all data}) d\phi_0 d\psi d \cos \iota$$

Preliminary results

Joint 95% upper limits from first ~ 13 months
of S5 using H1, H2 and L1 (97 pulsars)




Lowest h_0 upper limit:


PSR J1435-6100 ($\nu_{\text{gw}} = 214.0$ Hz, $r = 3.3$ kpc) $h_{0_min} = 4.2 \times 10^{-26}$

Lowest ellipticity upper limit:

PSR J2124-3358 ($\nu_{\text{gw}} = 405.6$ Hz, $r = 0.25$ kpc) $\varepsilon = 9.6 \times 10^{-8}$

Due to pulsar glitches the Crab pulsar result uses data up to the glitch on 23 Aug 2006, and the PSRJ0537-6910 result uses only three months of data between two glitches on 5th May and 4th Aug 2006





- **Black** curve represents one full year of data for all three interferometers running at design sensitivity
- **Blue** stars represent pulsars for which we are reasonably confident of having phase coherence with the signal model
- **Green** stars represent pulsars for which there is uncertainty about phase coherence

Crab pulsar – spin-down limit

- Spin-down limit assumes all the pulsars rotational energy loss is radiated by gravitational wave
- We know some goes is emitted electromagnetically and is powering the expansion of the Crab nebula
- This is poorly constrained and allows room for gravitational wave emission
- Braking index $\dot{\Omega} = -k \Omega^n, n = \frac{\nu \ddot{\nu}}{\dot{\nu}^2}$
 - » The braking index of the Crab is $n=2.5$, not $n=3$ for purely magnetic dipole radiation, and not $n=5$ for purely gravitational radiation emission
 - » Palomba (2000) allows for a combination of mechanisms to account for this braking index and ends up with a GW spin-down limit which is 2.5 times below the $n=5$ standard limit.



Standard spin-down limit:

$$h_0^{\text{spin-down}} = \left(\frac{5 G I_{zz} \dot{\nu}}{2 c^3 r^2 \nu} \right)^{1/2}$$



Crab pulsar - result

- These results give upper limits for the Crab pulsar of $\epsilon < 2.8 \times 10^{-4}$, $h_0 < 5.2 \times 10^{-25}$
 - » this value of the ellipticity is now in the range of some of the more speculative equations of state (Owen, 2005)
- These beat the spin-down limit of $h_0 < 1.4 \times 10^{-24}$ by a factor of 2.7 – for canonical moment of inertia $I = 10^{38} \text{ kgm}^2$ - we even beat Palomba's limit
- Start to constrain the amount of spin-down energy in GWs to less than 10% of overall emitted and known spin-down (Palomba, 2000, Santostasi).
 - » This is significant: the **uncertainties** on all non-GW contributions add up to 80% of the total!
- Moment of inertia is uncertain by about a factor of three, but we can plot the result on the moment of inertia – ellipticity plane to give exclusion regions (Pitkin for the LSC, 2005)



Multi-template search

- The time domain search only operates on a very narrow band – if the gravitational radiation time evolution is different from that of the electromagnetic radiation its possible it may be missing the gravitational waves
- Consider mechanisms by which any emitted gravitational waves will differ from the electromagnetic
- In general, we might assume the following relationship between the GWs and EWs: $f_{gw} = 2f_{ew}(1 + \delta)$ where $\delta \ll 1$
- In the case of free precession, where the spin axis is not aligned with the inertial axis: $\delta \sim \alpha \epsilon$ where $\alpha \sim 1$ and depends on the geometry and ϵ is the ellipticity
- Similarly, if we consider a two-component model, where the GW source spins down separately from the EW, but are connected by some torque acting over a coupling timescale:

$$\delta \sim \frac{\tau_{coupling}}{\tau_{spin-down}}$$

Search parameters continued

- In either of these cases, ϵ is approximately 10^{-4} . In one case roughly the largest imaginable ellipticity, and the other, $\tau_{coupling} \sim 10^{-4} \tau_{spindown} \sim 0.1 \text{ years}$ is roughly the longest timescale seen in glitch recovery
- Thus we are naturally led to a frequency band of $\Delta f_{gw} \sim 2f_{ew} \delta \sim 6 \times 10^{-3} \text{ Hz}$
- We also can work out $\Delta f_{gw}^{\cdot} \sim 3 f_{ew}^{\cdot} \delta \sim 1.2 \times 10^{-13}$ and $\Delta f_{gw}^{\cdot\cdot} \sim 8 f_{ew}^{\cdot} \delta \sim 1 \times 10^{-23}$

Conclusions

- Preliminary and expected upper limits for a large number of pulsars
- Have new timing for some pulsars – need more though over more of S5
 - » still need final timings and checks on parameter uncertainties
 - » produce final results after the end of S5
- Very interesting result for the Crab pulsar
 - » produce results of small parameter search
 - » Signal could be incoherent over glitch, so future analysis will use current result as prior on rest of S5 data
- Obtain more pulsar timing e.g. Terzan 5 pulsars and X-ray pulsar (PSRJ0537-6910)
- Carry on analysis...