





What was observed?













LIGO A signal from a binary black hole merger





LIGO A signal from a binary black LSC hole merger











The detector works



First-ever tests of Einstein's theories under dynamical, extreme-gravity conditions





Amazing Facts about GW150914



3 times the mass of the sun (1 million times the mass of Earth) turned into energy vibrating the fabric of spacetime

Peak power more than

Relative distance change same as **changing distance to nearest star by** width of human hair

Merger actually happened more than 1 billion years ago

In the volume that we can see systems like GW150914, there are more than **5 million galaxies**



LIGO Original detection strategy succeeds!



Proposal to the National Science Foundation

THE CONSTRUCTION, OPERATION, AND SUPPORTING RESEARCH AND DEVELOPMENT OF A

LASER INTERFEROMETER GRAVITATIONAL-WAVE **OBSERVATORY**

Submitted by the CALIFORNIA INSTITUTE OF TECHNOLOGY Copyright © 1989

Rochus E. Vogt Principal Investigator and Project Director California Institute of Technology

Ronald W. P. Drever Co-Investigator California Institute of Technology

Frederick J. Raab Co-Investigator California Institute of Technology Kip S. Thorne Co-Investigator California Institute of Technology

Rainer Weiss Co-Investigator Massachusetts Institute of Technology




Figure II-2 A comparison of the strengths of gravitational waves (characteristic amplitude h_c and frequency f) for burst signals from various sources (dashed lines and arrows), and benchmark sensitivities h_N (solid curves and stippled strips atop them) for interferometric detectors today (prototype) and in the proposed LIGO (early detector, advanced detector). See the caption of Figure A-4a (a duplicate of this figure) and the associated discussion in Appendix A for more details.

LIGO A signal from a binary black hole merger



